



# **B-tagging Performances**

***Simona Rolli***

***Tufts University***

***Third North American ATLAS Workshop***


***Boston, July 2006***

# Introduction



b-tagging is an important tool for physics at LHC:

- Aim is to tag inclusively high-pT b-jets
  - Searches:
    - Low mass SM Higgs (associate production with tt,W)
    - MSSM Higgs boson
    - SUSY
    - ...
  - Precision measurements
    - Top physics






# Outline of the talk

---

- Summary on taggers
- Performance
  - ♦ Different physics samples
  - ♦ Different releases
    - Tracking, generators, ...
- First Data and b-tagging commissioning
- Conclusions

# Building a tagger

- All taggers rely on a comparison between two hypotheses:
  - ♦ likelihood for the jet to come from a b-quark vs light quark
- Example: transverse impact parameter



- use the normalized impact parameter significance ( $S = d_0/\sigma_d$ ) of each track
- compare it to predefined calibration p.d.f. for the b and light hypothesis  
 ➔  $b(S)$  and  $u(S)$
- sum over all tracks ➔ jet btag weight

$$W_{\text{jet}} = \sum_{\text{tracks}} \ln \left( \frac{b(S)}{u(S)} \right)$$


- high  $W$  ➔ likely a b-jet

# Taggers available



	1 <sup>st</sup> stream (AOD and CBNT)	2 <sup>nd</sup> stream (AOD only)	
IP (long. impact)		Lifetime1D	needs good primary reconstruction
(trans. impact)	IP2D	Lifetime2D	the most robust
	IP3D	Lifetime3D	combination
Inclusive Secondary Vertex	SV1	SecVtxBU	more demanding for tracking
	SV2	SecVtxTD	
Pre-defined combination	“weight”: IP3D + SV1	“lhSig”	

# b-jet selection (Wt events)



From the BJet collection jets are selected using TruthInfo, LHSig and SV1



---

# Performances



# B-tagging performance estimators



- B-jet efficiency  $\varepsilon_b$  as function of variable cut:
  - ♦ Denominator:
    - jets defined as b using MC truth
      - with fixed  $p_T$  and  $\eta$  cuts ( $p_T > 50$  GeV/c,  $|\eta| < 2.5$ )
  - ♦ Numerator:
    - ditto + cut on a tagging weight
- Light-jet rejection  $R_u = 1 / \varepsilon_u$ 
  - ♦  $R=100$  means 1% mistag rate
  - ♦ light jets: u, d, s, g
- B-jet efficiency as a function of  $P_T$  and  $\eta$ 
  - ♦ Denominator:
    - jets defined as b using MC truth
      - with fixed cut on weight (SV1 > 3, LHSig > 0.9)
  - ♦ Numerator:
    - ditto + cut on  $p_T$  and  $\eta$





# Performance Studies

Many of the performance studies are motivated by physics studies

**It is important to study b-tagging for different samples and in the context of each physics analysis**

- The b-tagging performances depend (at least) on the jet  $p_T$ ,  $\eta$  and isolation (event topology)
- The optimization of b-tagging algorithms and the use of b-tagged jets in analysis cuts also depend on the specific physics analysis
- The understanding of b-tagging performances can be a significant systematics for the analyses which make use of this tool

- **SUSY**

Tommaso Lari (Milano)

- **Higgs**

WH validation -Marseille

WH validation (Siegen University)

Little Higgs (high  $p_T$  b-jets) (Valencia)

- **Top:**

Anne-Isabelle Etienne (Saclay)

Simona Rolli (Tufts)

Marisa Sandhoff (Wuppertal)

Marseille group

## Various software releases used

- Release Validation
- Tracking Validation
- Generators Validation (fragmentation, decays, etc)
- Jet Algorithms Dependence

# B-tag in SUSY Searches

- The most general signature of R-parity conserving Supersymmetry is **hard jets and missing energy**.
- Main backgrounds are **tt, W+jets, Z+jets, QCD multi-jets**
- b jets arises from gluino decays involving b and t squarks. They carry useful information on **third generation of scalar quarks** (enhanced if lighter than first two generations)
- b-tagging can be used to **reject SM backgrounds** (other than tt, bb+jets) and **discriminate between SUSY models/points** in parameter space.
- An analysis was made on Rome Full Simulation samples (tt, W+jets, SU1, SU2, SU3) to investigate the use of b-tagging in SUSY jet+missing Et inclusive searches:
  - ♦ W+jets background can be strongly suppressed by b-tagging ( $F_{abSV} > 4$  and  $P_T > 100 \text{ GeV}$ ).
  - ♦ Ongoing work on studying bbqq and Z+jets backgrounds.

	0 b-jets	1 b-jet	2 b-jets
tt	24	36	9
W+jets	307	48.6	0
SU1	122	57.1	15.3
S/\B	6.7	6.2	5.1
SU3	253	105	22
S/\B	13.9	11.4	7.4

Tommaso Lari, May 2006



# Performance in SUSY/Top events

## B-tagging as a tool to discriminate SUSY signal from background

- Analysis based on AOD produced with release 11.0.41 (official pre-CSC production) plus some older Rome samples (not yet available in CSC)
  - ♦ B-tagging using lhSig (CERN tagger) or weight (Marseille tagger)
  - ♦ Jets are cone 0.4, tracking with iPatRec
- Samples
  - ♦ **T1** (tt, MC@NLO, inclusive dileptonic+semileptonic)
  - ♦ **SU1, SU2, SU3, SU4** (Different SUSY points, JIMMY)
  - ♦ **A7** (W+jets, ALPGEN)
- Jet Flavor Identification
  - ♦ **b-jet**: b-quark within DR=0.3 cone
  - ♦ **c-jet**: c-quark within DR=0.3, no b-quark within DR=0.6
  - ♦  **$\tau$ -jet**: not identified by  $\tau$ -tagging, t within DR=0.3
  - ♦ **Light jet**: no b,c,  $\tau$  within DR=0.6



Some jets do not belong to any of above categories and are ignored

No isolation condition is used to select jets



Dependence on flavour tagging cone size weak. Basically same results if a  $\Delta R=0.8$  veto cone is used

Tommaso Lari, May 2006


# Performance in SUSY/Top events



Tommaso Lari, May 2006



# Performance in SUSY/Top events



- **b-tagging** (rejection of light jets) in **SUSY samples similar than in top events**, when rejection is plotted **as a function of jet pt**.
  - Exception: jets with very low pt ( $< 30$  GeV) poorly tagged in SUSY events. Need to be further studied
- It is important to **improve b-tagging for very high-pt jets**, which are present in many SUSY (and SUSY-like) events (see Valencia group studies).

Tommaso Lari, May 2006

# WH(120 GeV) benchmark 10.5.0

J.B.DeVivie




WH events,  $m_H = 120 \text{ GeV}/c^2$ ,  
(Rome layout, datasets 4860/4861)

b-jets:  $H \rightarrow b\bar{b}$

light jets:  $H \rightarrow u\bar{u}$

Release 10.5.0 + patches

xKalman tracks, Cone 0.4 jets



Rejection of light quark jets:

	IP2D	IP3D	SV1+IP3D	SV2+IP3D	lhSig (à la 10.5.0)
50% b-tag effi.	130	208	672	708	100
60% b-tag effi.	50	72	155	153	41

# WH(120 GeV) benchmark with 11.0.4x

WH events,  $m_H = 120 \text{ GeV}/c^2$ ,


Data sample 4860/1 :

**Rome** with 11.0.3 (private reco)

**mc11** with 11.0.41 (official AOD)

Release 11.0.3/4x ,  
iPatRec tracks, Cone 0.4 jets

Major change in the ATLAS layout  
between Rome and CSC : **three**  
**pixel layer geometry**



Clear improvement with the 3 layers,  
e.g. light jet rejection @  $\epsilon_b = 60$  (50) %

raw	IP2D	SV1
<b>Rome</b>	76 (270)	285 (1265)
<b>mc11</b>	99 (450)	324 (1550)

J.B.DeVivie, May 2006



# WH(120 GeV) benchmark with 11.0.4x: calibration functions



J.B.DeVivie, May 2006

Use different csc samples to build the calibration functions  
Samples 5200 (T1), 5340/1 (ttH), 5850/1/4/7 (WH120, 400)

⇒ Each jet/track enters the pdf with a weight 1

Predefined p.d.f. for b and  
light quark hypothesis

light jet rejection @  $\varepsilon_b = 60$  (50) %

	IP2D	IP3D	SV1+IP3D	SV2+IP3D
official AODs	75 (281)	109 (476)	192 (1024)	187 (1056)
with recalibration	74 (292)	111 (498)	232 (1222)	234 (1305)

Very small effect for IP (< 5%),  
Largest effect in SV (~ 20%) as  
expected (due in particular to the  
mean SVX finding efficiencies  
entering the weights :

(%)	light jets	b-jets
Rome	2.8	65.2
csc11	4.2	69.2

# WH(120 GeV) benchmark with 11.0.4x: Parton Shower



J.B.DeVivie, May 2006

Data sample

- 4860/1 : **mc11** → old shower (PYEVNT)
- 5850/1 : **csc11** → new shower (PYEVNW)


Light jet rejection as a function of  $|\eta|$   
and  $p_T$  @  $\epsilon_b = 60\%$  , for **IP2D**

very large degradation, back to Rome  
performances (or worse)  
with a 3 pixel layer layout...


very annoying effect since we do not  
really know which shower model is  
closest to reality... (PYEVNW should be  
better but is it properly tuned ?)

**IP2D** and **SV1** @  $\epsilon_b = 60$  (50) %

raw	<b>IP2D</b>	<b>SV1</b>
<b>mc11</b>	<b>99 (450)</b>	<b>324 (1550)</b>
<b>csc11</b>	<b>75 (281)</b>	<b>192 (1024)</b>



Mislabelling of b-quark in PS  
Corrected in 12.0.x



# WH benchmark with 11.5.0: Tracking Algs

WH events generated with 11.0.41 (CSC sample A)  
Reconstruction 11.5.0  
DC3-02 layout

Two taggers considered for performance studies:

Marseille

CERN

Two tracking algorithms used:



xKalman

newTracking

V. Sipica, Siegen May 2006

Particle jets with cone 0.4

Jet weights computed with 11.5.0



# WH benchmark with 11.5.0: Tracking Algs

V. Sipica, May 2006

- Performances of Marseille taggers in different steps:

- 'just taggers' => useSharedHits & rejectBadTracks set to false

- 'V0 rejection' => rejectBadTracks set to true

- Standard cuts are used to calculate the rejection

**xKalman**

11.5.0	IP2D		IP3D		IP3D + SV1	
Efficiency	50%	60%	50%	60%	50%	60%
just taggers	175 +/- 8	62 +/- 2	301 +/- 18	98 +/- 3	690 +/- 64	172 +/- 8
+V0 rejection	277 +/- 16	77 +/- 2	432 +/- 32	115 +/- 4	816 +/- 82	192 +/- 9

- Standard cuts are used to calculate the rejection

**newTracking**

11.5.0	IP2D		IP3D		IP3D + SV1	
Efficiency	50%	60%	50%	60%	50%	60%
just taggers	230 +/- 12	80 +/- 2	528 +/- 42	137 +/- 6	937 +/- 100	255 +/- 14
+V0 rejection	393 +/- 27	107 +/- 4	654 +/- 58	161 +/- 7	1055 +/- 119	267 +/- 15

- newTracking / xKalman:

11.5.0	IP2D		IP3D		IP3D + SV1	
Efficiency	50%	60%	50%	60%	50%	60%
just taggers	1.32 +/- 0.07	1.28 +/- 0.04	1.76 +/- 0.10	1.39 +/- 0.06	1.36 +/- 0.14	1.48 +/- 0.07
+V0 rejection	1.42 +/- 0.09	1.39 +/- 0.05	1.51 +/- 0.12	1.4 +/- 0.06	1.29 +/- 0.15	1.39 +/- 0.07

Similar improvement for CERN taggers

# b-tagging in $t\bar{t}$ events: 10.0.1

- $t\bar{t}$  MC@NLO+Jimmy (sample 4100 T1)
- From AODs, cone  $\Delta R=0.4$ , iPatrec tracks
- b and light jets from same sample: purification
- Statistics: 395k b-jets, 618k(517k) u-jets

Release 10.0.1 + patches

L. Vacavant, Mar 2006

	$R_u (\epsilon_b=50\%)$ (raw)	$R_u (\epsilon_b=60\%)$ (raw)	$R_u (\epsilon_b=50\%)$ (purified)	$R_u (\epsilon_b=60\%)$ (purified)
IP2D	<b><math>158 \pm 3</math></b>	<b><math>55 \pm 1</math></b>	<b><math>181 \pm 3</math></b>	<b><math>60 \pm 1</math></b>
IP3D	$228 \pm 4$	$86 \pm 1$	$258 \pm 6$	$94 \pm 1$
SV1+IP3D	<b><math>505 \pm 14</math></b>	<b><math>184 \pm 3</math></b>	<b><math>773 \pm 30</math></b>	<b><math>240 \pm 5</math></b>
Lifetime2D	$145 \pm 2$	$53 \pm 1$	$169 \pm 3$	$57 \pm 1$
lhSig	$172 \pm 3$	$66 \pm 1$	$200 \pm 4$	$73 \pm 1$

More in A.I.Etienne studies in the Top Group (Effects of environment)

# b-tagging in ttbar: 11.5.0 vs 12.0.0



Reco	12.0.0	11.5.0	F(12/11)
2D 60%	$84 \pm 2$	$75 \pm 4$	$1.12 \pm .06$
3D 60%	$133 \pm 4$	$130 \pm 10$	$1.02 \pm .08$
3DSV1 60%	$296 \pm 12$	$319 \pm 37$	$0.93 \pm .11$

CSC tt sample 5200,  
Simulation 11.0.4,  
Reconstruction 11.5.0/12.0.0,  
VxPrimary, no shared hits,  
b-jet efficiency 60%, iPatrec,  
old Rome calibration

CSC tt sample 5200,  
Simulation 11.0.4,  
Reconstruction 11.5.0/12.0.0,  
VxPrimary, no shared hits,  
b-jet eff 60%, newTracking,  
old Rome iPat calibration

Reco	12.0.0	11.5.0	F(12/11)
2D 60%	$81 \pm 5$	$92 \pm 6$	$0.88 \pm .07$
3D 60%	$108 \pm 8$	$154 \pm 13$	$0.70 \pm .08$
3DSV1 60%	$171 \pm 15$	$265 \pm 28$	$0.65 \pm .08$

**Software issues to be checked: clustering, calibration DB, multiple collections treatment etc**

A. Rozanov, , June 2006



# b-tagging in Single Top: 10.0.1




## Decay modes:

- $W^*$  :  $W^* \rightarrow t \bar{b} \rightarrow (l^+ \nu_b) \bar{b}$
- $Wg$ :  $q'g \rightarrow t q \bar{b} \rightarrow (l^+ \nu_b) q \bar{b}$
- $W+t$ :  $bg \rightarrow t W \rightarrow (l^+ \nu_b) qq'$


1 leptons + MET  
+  $\geq 2$  jets  
+ 1(2) b-tags





## Release 10.0.1 + patches



## W\* (s-channel)



S. Rolli, June 2006





# Wt: b-tagging efficiencies

In respect to tagger cut

Denominator: number of jets matched with the b-parton (Tinfo), with  $P_T > 50$  GeV,  $\eta < 2.5$

Numerator: ditto with cut on weight/likelihood

IP2D Cut	Eff Ip2D	SV1 Cut	Eff SV1	LHSig cut	Eff LHSig
1	0.60 0.63	1	0.63 0.63	0.1	0.80 0.75
2	0.54 0.55	2	0.59 0.59	0.2	0.76 0.72
3	0.49 0.48	3	0.55 0.57	0.3	0.72 0.69
4	0.43 0.41	4	0.53 0.54	0.4	0.70 0.67
5	0.38 0.35	5	0.51 0.51	0.5	0.68 0.66
6	0.33 0.28	6	0.48 0.48	0.6	0.67 0.65
7	0.29 0.21	7	0.46 0.46	0.7	0.65 0.63
8	0.25 0.18	8	0.43 0.43	0.8	0.63 0.61
9	0.21 0.14	9	0.41 0.40	0.9	0.60 0.57

Cone 0.7



Cone 0.4

ATL-PHYS-COM-036

In respect to PT and  $\eta$

Denominator: number of b-partons with  $P_T$  and  $\eta$  in given interval;

Numerator: bjets matched with the b-parton (parton level info) with  $P_T$  and  $\eta$  in given interval and cut on weight/LHSig.



# Remarks on Current Taggers


- All taggers are kept for performance studies and cross-checks
  - ⇒ **low performance taggers** (Lifetime2D/IP2D) are **usually rather robust** (easier to understand and commission)
  - ⇒ **high performance** ones (SV1/SV2) will **require more time** to control
  - ⇒ taggers identical wrt discriminating variables (Lifetime2D ~ IP2D, Lifetime3D ~ IP3D) are kept for cross-checks and do differ in some point (refined track selection in IPxD, one 2D vs one 1D pdf for IP3D vs Lifetime3D, ...)
- For physics analysis, a combination is given :
  - 🏷️ **“1<sup>st</sup> stream” taggers** : (\*JetTag)→weight()  
corresponding to SV1+IP3D  
**← most powerful tagger**
  - 🏷️ **“2<sup>nd</sup> stream” taggers** : (\*JetTag)→weightForTag(“lhSig”)  
corresponding to Lifetime1D+Lifetime2D(+SecVtxBU)

# Soft Lepton Taggers: Muon


## Soft Muon Tagger

- Implementation in the JetTagging package finished (SoftMuonTag).
- Selection:
  - STACO or MuTag muon
  - $\Delta R < \Delta R^{cut}$  (default: 0.6). Jet momentum corrected for the presence of the muon (i.e. muon momentum added to jet momentum).
  - $p_T > p_T^{cut}$  (default: 4 GeV)
  - $|d_0| < d_0^{cut}$  (default: 4 mm) 2D impact parameter w.r.t. primary vertex.

Efficiency (Muons from B), Barrel



$p > 3\text{GeV}$



# Soft Lepton Taggers:Electron





## Soft Electron Tagger

Basic soft electron btagging tools implemented

- ECAL +TRT (+Si)

For each track in the jet:

- soft electron weight from ElectronCollection
- max electron weight stored in SoftLeptonInfo





# First Data and b-tagging Commissioning



# Ongoing Activities

---

- Calibration Issues - see previous talk
- Tools for Validation
- Commissioning btag with top data

# Tools for Validation

Standard Set of Tools to compare the performances from various algorithms


A set of standalone (ROOT only) classes to produce

- 1) Histo and plots of various variables
- 2) 2D Histos/plots rejection vs b-efficiency
- 3) mis-identification for a fixed b-efficiency

Freiburg University

Histograms can be used to create pdf (likelihood ratio)

The Root tree can be used to train a NN for combination



**Work in progress to  
integrate with current  
releases  
(now based on 10.0.4)**





# Lifetime Efficiency from top data

It is necessary to evaluate the taggers performance from data

At the Tevatron this was done in two ways:

H. Bachacou, May 2006

- rejection from jet samples (previous talk)
- b-tag efficiency from semileptonic decay and cross calibration with soft lepton/lifetime taggers

At LHC the high statistics  $t\bar{t}$  sample can be used

- relatively pure b-jet sample
- inclusive b-decay (not semileptonic only)
- high  $E_T$  sample
- large heavy flavor content:
  - 2 b-jets/events
  - 1 c-jet/2 events from  $W \rightarrow jj$
- it assumes  $\text{Br}(\text{top} \rightarrow Wb) = 100\%$

Method: count the number of events with 1,2,3 tagged jets and extract  $\epsilon(b), \epsilon(c), \sigma(t\bar{t})$  using a likelihood fit


# Lifetime Efficiency from top data



Several pseudo-experiments ( $\sim 15 \text{ pb}^{-1}$  each)

H. Bachacou, May 2006

	'True'	Measured
$\epsilon_B$ (%)	$67.9 \pm 0.2$	$66.8 \pm 2.3$
$\epsilon_C$ (%)	$20.3 \pm 0.4$	$28.5 \pm 5.3$
$\epsilon_L$ (%)	$0.93 \pm 0.03$	-
$\sigma_{t\bar{t}}$ (pb)	857	$878 \pm 28$





Working on systematics and improving event selection

Background evaluation needs more refinement (W+jets ok, QCD?)

- topological likelihood discriminant ?

Dilepton channel can be also used (no c-jet component)

# B-tagging commissioning

- Already lots of activities ongoing
  - ♦ Many more will start and many more will be repeated when first data will be available
    - It's important to have the machinery ready
  - ♦ Many variables to test.... 
    - Detector commissioning (for example effects of misalignment)
    - Simulation issues
      - Tracking, material....
  - ♦ not for the first time... 
    - TeVatron experience!
  - ♦ Much more data than at the TeVatron
    - Top sample:
      - « pure » b sample:  $t\bar{t}$  semi-leptonic (and dilepton)
  - ♦ « pure » light quark sample (very high purity needed): W/Z+jet ? di-jets ?
  - ♦ « pure » c sample: ?

# B-tagging and first physics results




- Top
  - ♦ will be used for commissioning
    - b-tag will be an ingredient
    - known physics process, useful for testing performance

- Searches

- ♦ “easy” searches don’t need b-tagging immediately
  - $Z', LQ, ED$ , mass bumps...
  - Third generation searches needs it...
    - cross check with Top

- Higgs

- ♦ Benchmark for performance of the ATLAS detector
  - Energy resolution,  $E_{\text{miss}}$ , lepton/photon ID
- ♦ Several of the prominent channels do not make use of b-tagging
  - 4 leptons,  $\gamma\gamma$
  - $t\bar{t}H$  does



# Conclusions

---

- Many performance tests ongoing
  - ◆ Driven by physics studies
    - From DC1 to CSC
  - ◆ Driven by performance tests
    - Tracking
    - Releases
- Work starting up on b-tagging calibration and performance validation with first data
  - ◆ Top sample



# Backup Slides

# Labelling and purification

---

- **Labelling:**
  - ♦ To quote performances we need to know the « true » flavour of the jet: not completely trivial...
  - ♦ Method used here: label a jet as a b-jet if there is a b-quark (after FSR\*) within  $\Delta R < 0.3$ . If not look for a c-quark  $\rightarrow$  c-jet. Otherwise jet is labelled as a light jet.
- **Overlapping jets and purification:**
  - ♦ Overlaps in jetty events  $\rightarrow$  mislabelling
  - ♦ Jet isolation very dependent on the type of events and physics processes (gluon jets) + jet algorithm
  - ♦ Purification may be used to factorize this from pure b-tagging issues: do not consider lights jets where there is a b/c/quark/hadron within  $\Delta R < 0.8$




# B-tag efficiencies




Cross check of LHSig distribution using a different tagger as selector.

LHSig distribution:  
IP2D > 3.0 (red)  
IP2D < 1.0 (black)



# High PT B-jets (in LH models)

ATL-PHYS-PUB-2006-003



$p_T$  (b-jets)  $\sim$  250 GeV


$Z_H \rightarrow$   $\left\{ \begin{array}{ll} b \bar{b} & 20000 \text{ events} \\ u \bar{u} & 20000 \text{ events} \\ c \bar{c} & 20000 \text{ events} \end{array} \right.$

ATHENA release 10.0.1 - Full Simulation

Room for further development


$p_T$  ( $b_1$ )  $\sim$  250 GeV

$p_T$  ( $b_2$ )  $\sim$  500 GeV





# b-tagging in ttbar events: 10.0.1

Rome sample 4100,  
semileptonic ttbar events  
Reconstruction 10.0.1  
b-tag rerun with patches



— Light jets  
— b jets  
— c jets

lhSig




	$R_1 (\epsilon = 50 \%)$	$R_1 (\epsilon = 60 \%)$
SV1 + IP3D	$593 \pm 8$	$207 \pm 3$
lhSig	$132 \pm 4$	$53 \pm 2$

A.I. Etievre, Dec 2005

# b-tagging in $t\bar{t}$ events: 10.0.1

## Environment

Comparison of the b-tagging efficiency in the two sides (leptonic/hadronic)



→ Nearly identical !

For a b-weight cut = 3:

	Global	Leptonic environment	Hadronic environment
Efficiency (%)	$64.5 \pm 0.1$	$64.6 \pm 0.1$	$64.3 \pm 0.1$

A.I. Etienne, Dec 2005



# b-tagging in Top Physics

---

- **In ttbar events, b-tagging of a jet is defined as  $SV1 + IP3D \geq 3$**   
(SV2 + IP3D when available)
- **b-tagging performances obtained with this cut on cone 0.4 size jets:**
  - Light jet rejection =  $116 \pm 4$
  - b-tagging efficiency =  $64.5 \pm 0.1 \%$
  - c jet rejection =  $6.1 \pm 0.1$
- **b-purity in top mass measurement:**  
 $63.2 \pm 0.9 \%$
- **For top mass measurement, these performances are perfect, but any advice in order to optimize the b-tagging performances is welcome!**

A.I. Etiennev, Dec 2005


# ttH vs ttjj : 10.0.1

- The most complicated (interesting!) of the benchmarks
  - ♦ busy events with overlaps, mislabelling issues
- b-jets: ttH Pythia (samples 4867, 4868) 20k evts
- u-jets: tt(jj) MC@NLO
  - ♦ tt (sample 4100 T1) 243k evts
  - ♦ tt(jj) (samples 4870,4871) 259k evts
    - Filter to select high jet multiplicity (j6pT14)
    - $\epsilon = 44\%$
- From AODs, cone  $\Delta R=0.4$ , iPatrec tracks, Release 10.0.1 + patches
- Statistics: 75k b-jets, 1.2M u-jets (Caveat: not the same generator)

L. Vacavant, Mar 2006

	$R_u$ ( $\epsilon_b=50\%$ )	$R_u$ ( $\epsilon_b=60\%$ )	$R_u$ ( $\epsilon_b=70\%$ )
<b>IP2D</b>	<b><math>218 \pm 3</math></b>	<b><math>66 \pm 1</math></b>	<b>23</b>
<b>SV1+IP3D</b>	<b><math>882 \pm 24</math></b>	<b><math>297 \pm 5</math></b>	<b>59</b>

# b-tagging in $t\bar{t}b\bar{b}$ events: 11.0.41



MC@NLO+HERWIG (10000  $\sqrt{s}$  evts)

	e=50%	e=60%
$R_e$	10 / 13	6 / 8
$R_{uds}$	500 / 680	90 / 230

→ Results improved compared to Rome by 20% for  $R_e$  and 50% for  $R_{uds}$  at e=60%

→ Need to check with the background when simulated.

P. Pralavorio, , March 2006



# Digression on Tracking eff (11.5.0)



## Integrated Tracking eff and fake rates (basic selection)



	11.0.41 Eff (%)	11.0.41 Fake (%)	11.5.0 Eff (%)	11.5.0 Fake (%)
IP	97.3	4.02	99.6	0.47
XK	97.7	2.08	99.3	0.54
NT	95.7	5.46	99.3	0.34

Cumulative Efficiency: 99.87%  
Sorting tracks and removing duplicates  
(IP: 99.6%, XK/NT: 99.3%)



IP  
XK  
NT

Venn Diagram for overlap





# Tracking Eff comparisons

CSC tt sample 5200, simu 11.0.4, reco 11.5.0, VxPrimary, no shared hits, no cleaning, small stat. 10K (updated 31.05.06)

Track finding	xKalman	xKalman	iPatrec	iPatrec	newTrack	newTrack
Calibration	Old xKalman	Old iPatrec	Old iPatrec	Old xKalman	Old newTrack	Old iPatrec
2D 60%	$70 \pm 4$	$70 \pm 4$	$75 \pm 4$	$71 \pm 4$	$70 \pm 4$	$92 \pm 6$
3D 60%	$108 \pm 7$	$104 \pm 7$	$130 \pm 10$	$137 \pm 10$	$108 \pm 7$	$154 \pm 13$
3DSV1 60%	$189 \pm 17$	$195 \pm 18$	$319 \pm 37$	$316 \pm 37$	$218 \pm 21$	$265 \pm 28$

Better numbers with cleaning



A. Rozanov, June 2006

# b-tagging and Jet algorithms

“jump” at efficiency of  $\sim 0.8$  from peak of weights for light jets  
Cone 0.7 and kT have similar behaviour

80k ttbar events, DC3, 11.0.41

CERN efficiency vs light jet rejection



Marisa Sandhoff

# Wt: Efficiencies ( $P_T$ and $\eta$ )





Efficiencies are calculated in the following way:

Denominator: number of b-partons with  $P_T$  and  $\eta$  in given interval;

weight/likelihood  
cut fixed

Numerator: bjets matched with the b-parton (parton level info) with  $P_T$  and  $\eta$  in given interval and cut on weight/LHSig.

S. Rolli, June 2006



# b-tagging in Single Top

## • B-Tag studies on $Wt$ , $W$ -g, $W^*$ AOD samples:

- Preliminary tests on various b-tag algorithms, as out of the box on Rome samples for single top were performed
- Reprocessing of data to obtain cone 0.4 bjets was done;
- Generally good agreement with other samples studies
- LHSig has slightly higher efficiency to select b-jets  
(LHSig > 0.9) in the data but has a very poor rejection factor.
- SV1 has slightly lower efficiency, but much higher rejection factor.

<b><math>Wt</math></b>	$R_u (\epsilon_b = 50\%)$	$R_u (\epsilon_b = 60\%)$
IP2D	<u>166</u> (125) (158-109)	<u>25</u> (50) (55-57)
LHSig	<u>NA</u> (172-NA)	<u>33</u> (33) (66-NA)
SV1 + IP2D	<u>333</u> (100) (505-325)	<u>100</u> (33) (184-156)

S. Rolli, March 2006

# Impact of (mis)alignment

Use CDF experience:

S. Gibson, Oxford

Map CDF commissioning misalignments from CDF run II to ATLAS and propagate to **b-tagging** performances (via degradation in track momentum, d0 resolutions)


		<i>2D algorithm</i>		<i>3D algorithm</i>	
		Perfect	Misaligned	Perfect	Misaligned
$R_u(\epsilon_b)$	$\epsilon_b = 50\%$	$155 \pm 8$	$138 \pm 6$	$348 \pm 22$	$316 \pm 19$
	$\epsilon_b = 60\%$	$47.1 \pm 1.1$	$43.6 \pm 1.0$	$89.0 \pm 3.4$	$81.7 \pm 2.7$
$\frac{R_u^{\text{Misaligned}}}{R_u^{\text{Perfect}}}$	$\epsilon_b = 50\%$	0.89		0.91	
	$\epsilon_b = 60\%$	0.92		0.92	

→ ~10% degradation  
→ Equivalent to 10  $\mu\text{m}$  r $\phi$  misalignment

Random misalignments:

IP3D tagger, ttH events, realistic conditions  
(redo exercise with detector misaligned from simu)

improving alignment up to  
almost perfect in one year,  
→ recovery of light jet rejection



# Lifetime Efficiency from top data

Working hypotheses:

- $\text{Br}(\text{top} \rightarrow \text{Wb}) = 100\%$
- $\epsilon(\text{b}), \epsilon(\text{c}), \epsilon(\text{light})$  assumed independent of number of tags

H. Bachacou, May 2006

Inputs:

- backgrounds need to be evaluated for each sub-sample
- $\epsilon(\text{light})$  needs to be measured somewhere else (for now simulation,  $\epsilon(\text{light}) = 0.01$ )
- $\sigma(\text{tt})$  measurement ingredients: trigger eff, ID eff, acceptance etc...

- $L = \text{Poisson}(N_1, \langle N_1 \rangle) \cdot \text{Poisson}(N_2, \langle N_2 \rangle) \cdot \text{Poisson}(N_3, \langle N_3 \rangle)$

- $N_n$  = Number of observed events with n tags

- $\langle N_n \rangle$  = Expected number of events with n tags : function of  $\epsilon_B, \epsilon_C, \epsilon_L, \sigma_{t\bar{t}}$ , etc...

$$\langle N_n \rangle = (L \cdot \sigma_{t\bar{t}} \cdot A_{\text{pre-tag}}) \cdot \sum_{i,j,k} F_{i,j,k} \cdot \lambda_{\text{combi}} \cdot A_i^{i'} \cdot \epsilon_b^{i'} \cdot (1 - \epsilon_b)^{i-i'} \cdot \epsilon_c^{j'} \cdot (1 - \epsilon_c)^{j-j'} \cdot \epsilon_l^{k'} \cdot (1 - \epsilon_l)^{k-k'}$$

$$i = \# \text{ b-jets and } i' = \# \text{ tagged b-jets}$$

$$j = \# \text{ c-jets and } j' = \# \text{ tagged c-jets}$$

$$k = \# \text{ l-jets and } k' = \# \text{ tagged l-jets}$$

$$F_{i,j,k} = \text{Fraction of events with } i \text{ b-jets, } j \text{ c-jets, } k \text{ l-jets.}$$

$$A_i^{i'} = i! / (i'! \cdot (i - i')!)$$

$$\sigma_{t\bar{t}} = \text{production cross-section}$$

$$A_{\text{pre-tag}} = \text{acceptance without b-tagging}$$

$$L = \text{integrated luminosity}$$